

Enantioselective Halocyclization Using Reagents Tailored for Chiral Anion Phase-Transfer Catalysis

Yi-Ming Wang, Jeffrey Wu, Christina Hoong, Vivek Rauniyar, and F. Dean Toste*

Department of Chemistry, University of California, Berkeley, California 94720, United States


S Supporting Information

ABSTRACT: A chiral anion phase-transfer system for enantioselective halogenation is described. Highly insoluble, ionic reagents were developed as electrophilic bromine and iodine sources, and application of this system to *o*-anilidostyrenes afforded halogenated 4*H*-3,1-benzoxazines with excellent yield and enantioselectivity.

The enantioselective synthesis of halogenated compounds is an important target for the synthetic community.¹ In particular, electrophilic halogenation of alkenes offers direct access to bifunctionalized molecules with the concomitant formation of up to two stereocenters. There are, however, only a limited number of approaches for the induction of enantioselectivity in this class of electrophilic addition reactions, especially in a catalytic sense.² In addition to transition metal-catalyzed alkene halogenations,³ organocatalytic methods have recently been reported using modified cinchona alkaloids,⁴ aminourea⁵ and bis(amidine) derivatives,⁶ C₃-symmetric imidazoline derivatives,⁷ and phosphoric acids⁸ as catalysts. While high enantioselectivities can be achieved, low temperatures and/or high catalyst loading are often required to overcome the uncatalyzed background reaction.

Very recently, our laboratories reported an enantioselective fluorocyclization of olefins using *chiral anion phase-transfer catalysis*,⁹ (Scheme 1) by taking advantage of the low solubility

Scheme 1. Application of Chiral Anion Phase Transfer Catalysis to Bromo- And Iodocyclization^a



^aPhos*⁻ = chiral phosphate anion, E = Br or I.

of the cationic fluorinating reagent Selectfluor in nonpolar solvents. Exchange of the tetrafluoroborate anions from the reagent with lipophilic chiral phosphate anions brings the active electrophile into solution as a chiral ion pair, resulting in efficient, chemo- and enantioselective fluorination of the substrate at or near room temperature. Notably, due to the

insoluble nature of the fluorinating reagent, nonselective background reaction is minimal. We hypothesized that other electrophilic functionalization reactions should be amenable to chiral anion phase-transfer catalysis, thereby extending the applicability of the concept to a general approach for catalysis. Toward this goal, novel electrophilic reagents needed to be developed where insolubility, normally detrimental, is a required feature. We report herein the synthesis of a new brominating reagent (**1d**) for use in anionic phase-transfer bromination, as well as an analogous reagent (**5d**) for iodination.

Given the lack of methods for enantioselective synthesis of 4*H*-3,1-benzoxazines,¹⁰ we chose the 6-*exo*-trig cyclization of amide **3a** to examine the utility of the cationic brominating reagents in chiral anion phase-transfer catalysis. (Bis)amine-halonium reagents have long been known and employed as electrophilic halonium sources in reactions with alkenes;¹¹ however, attempts to use the known oligomeric DABCO-bromine complex or bis(*sym*-collidine)bromine(I) hexafluorophosphate under chiral anion phase-transfer conditions resulted in rapid conversion to **4a** with poor enantioselectivity, reflecting a significant amount of uncatalyzed background reaction (Table 1, entries 1 and 2).¹²


The lack of success using these monocationic reagents led us to consider the monoalkylated-DABCO scaffold as a starting point, in analogy to Selectfluor, for the generation of more highly charged reagents for phase-transfer bromination. To that end, treatment of monoalkyl DABCO tetrafluoroborates with elemental bromine in the presence of AgBF₄ resulted in the formation of tricationic brominating reagents **1**, which could be isolated by precipitation (Figure 1). These brominating reagents could be stored in an airtight vial at -20 °C for at least a month without deterioration. Moreover, **1a** was found to mediate the bromocyclization of amide **3a** almost instantaneously under homogeneous conditions in acetonitrile. In contrast, **1a** was insoluble in organic solvents of low or moderate polarity, including hexanes, toluene, and dichloromethane, and exposure of amide **3a** to **1a** in toluene did not result in significant formation of the cyclized product. Encouragingly, when **3a** was combined with **1a** in the presence of chiral phosphoric acid **2a** (10 mol %) and Na₂CO₃, desired product was obtained in moderate enantioselectivity (Table 1, entry 3).

With these promising results, we sought to optimize the structure of the brominating reagent. To increase the phase-

Received: June 14, 2012

Published: July 25, 2012

Table 1. Optimization of Enantioselective Bromocyclization



entry	"Br ⁺ "	cat.	solv	base	time	conv. ^a	ee ^b
1	DABCO·2Br ₂	2a	toluene	Na ₂ CO ₃	0.5 h	100%	39%
2	(coll) ₂ Br ⁺ PF ₆ ⁻	2a	toluene	Na ₂ CO ₃	0.5 h	100%	1%
3	1a	2a	toluene	Na ₂ CO ₃	24 h	82%	59%
4	1b	2a	toluene	Na ₂ CO ₃	24 h	67%	70%
5	1c	2a	toluene	Na ₂ CO ₃	24 h	30%	20%
6	1d	2a	toluene	Na ₂ CO ₃	24 h	67%	83%
7	1d	2a	<i>p</i> -xylene	Na ₂ CO ₃	16 h	62%	86%
8	1d	2a	xyl/hex (1:1) ^c	Na ₂ CO ₃	16 h	90%	89%
9	1d	2a	hexanes	Na ₂ CO ₃	16 h	21%	60%
10	1d	2b	xyl/hex (1:1) ^c	Na ₂ CO ₃	17 h	90%	89%
11	1d	2c	xyl/hex (1:1) ^c	Na ₂ CO ₃	17 h	90%	91%
12	1d	2c	xyl/hex (1:1) ^c	Na ₂ CO ₃ ^d	17 h	100%	90%
13	1d	2c	xyl/hex (1:1) ^c	Na ₃ PO ₄ ^d	17 h	100%	92%
14	1d ^e	2c ^f	xyl/hex (1:1) ^{c,g}	Na ₃ PO ₄ ^d	4 h	100% ^h	94%

^aEstimated from ¹H NMR of the crude product. ^bDetermined by chiral HPLC. ^c*p*-Xylene and hexanes. ^dFinely ground and dried overnight. ^eReprecipitated from MeNO₂. ^f5 mol %. ^g0.025 M. ^h82% isolated yield.




Figure 1. Synthesis of cationic halogenating reagents and ORTEP diagram of **1d** (ellipsoids at 50% probability level, counterions and H's omitted for clarity).


transfer efficiency and electrophilicity of the brominating reagent, we synthesized the more lipophilic and electron-deficient reagent **1b**. Indeed, treatment of **3a** with **1b** resulted in product formation in improved enantioselectivity. While a further increase in electron-deficiency (**1c**) was detrimental, the use of more lipophilic reagent **1d** afforded **4a** with increase in enantioselectivity to 83% ee (Table 1, entries 3–6). A decrease in the polarity of the solvent to 1:1 *p*-xylene/hexanes resulted in better conversion and enantioselectivity (entries 7, 8). While commercially available TRIP (**2b**) catalyzed the reaction with nearly identical efficiency and selectivity, a more highly lipophilic phosphoric acid with 6,6'-TIPS substituents (**2c**, TIPS-TRIP) resulted in further improvements (entries 10, 11). Use of **1d** reprecipitated from nitromethane (see Supporting Information) shortened reaction time without affecting enantioselectivity, and optimization of base and concentration improved enantioselectivity to 94% ee in 82% isolated yield

with no loss in enantioselectivity observed when catalyst loading was decreased to 5 mol % (entry 12–14).

With optimized reaction conditions in hand, we explored the substrate scope of our bromination reaction (Table 2). We were pleased to find good tolerance for substituents of varying steric bulk at the α position of the styrenyl system. Potentially problematic functional groups including a silyl ether (**4e**), an unprotected alcohol (**4f**), and a tertiary amine (**4o**) did not significantly impact the reactions. Notably, a substrate with a pyridyl backbone gave desired product in excellent yield and enantioselectivity (**4n**). Electron-withdrawing (**4q**) and electron-donating (**4r**) substitution at the *para* position, as well as *ortho* substitution (**4p**) of the benzamide aryl were tolerated. A slight modification of reaction conditions also afforded benzoxazine **4s** with *t*-butyl substitution at the 2-position. Gratifyingly, heteroarylamides (**3t–3v**) were also suitable substrates under standard reaction conditions. These examples, all run at room temperature without precaution to exclude air and moisture, demonstrate the mildness and operational simplicity of the reaction conditions.

We next explored substrates bearing substitution at both the α and the β positions of the styrenyl system. In the event, subjection of these substrates to standard reaction conditions furnished desired products in excellent yields and enantioselectivities. In all cases, only one diastereomer was observed in the ¹H NMR of the crude product. Notably, substituents of varying steric and electronic properties were tolerated (**4g–4i**). For example, enantioenriched *gem*-bromochloro adduct **4h** was formed in 88% yield and 96% ee from the corresponding vinyl chloride. Cyclic substrates, including conjugated (**4k**) and

Table 2. Scope of Enantioselective Bromocyclization




^aConditions: **1d** (1.3 equiv), TIPS-TRIP (5 mol %), Na₃PO₄ (4.0 equiv), *p*-xylene/hexanes (1:1, 0.025 M), 4 h, rt. Relative and absolute stereochemistry assigned by analogy to that of **4l**, determined by X-ray crystallography. Yields after chromatographic purification and enantioselectivities determined by chiral HPLC. ^b1.5 equiv **1d** in mesitylene/hexanes (1:1).

remote dienes (**4l**), afforded desired product with excellent chemo- and enantioselectivity. Similarly, benzothiophene substituted substrate **3m** provided dearomatized product in excellent yield and enantioselectivity. These results highlight the extraordinary tolerance of this catalyst system for substitution on the alkene moiety.


We speculated that enantioselective iodination might also be achieved through an analogous reagent. Thus, we prepared iodination reagent **5d** as a further test of the robustness and generality of the anionic phase-transfer approach. Through a procedure nearly identical to the one used in the preparation of

1d, reagent **5d** was obtained as an off-white solid of similar stability. We were pleased to find that treatment of **3b**, **3j**, and **3n** with **5d** (in place of **1d**) for 16 h without otherwise modifying reaction conditions furnished the corresponding iodinated products **6b**, **6j**, and **6n** in similarly high yields and enantioselectivities (Scheme 2).

Scheme 2. Enantioselective Electrophilic Iodocyclization



To provide some evidence that the reaction is proceeding through the proposed anionic phase-transfer process, we subjected **3a** to optimized reaction conditions but in the absence of base, which is required for the formation and regeneration of the chiral phosphate anion. After 2 h reaction time, the product was obtained in moderate conversion. The enantioselectivity observed, however, was greatly diminished (eq 1). On the other hand, use of base alone without **2c** did not



With 4.0 equiv Na₃PO₄:
No Base:

>98% conv, 94% ee
40% conv, 49% ee

result in rate acceleration. Thus, the observed rate and enantioselectivity enhancement in the presence of base cannot be attributed to deprotonation of the amide substrate. These experiments strongly suggest that the chiral phosphate anion is the active catalytic species in our optimized system (and not the phosphoric acid). Moreover, a nonlinear effect study (see Supporting Information) showed significant deviation from linearity. This study, combined with the observation that monoalkyl DABCO-derived reagents provide products with increased enantioselectivity, suggests that the occurrence of highly charged intermediates may be advantageous for enantioinduction in chiral phosphate phase-transfer catalysis.

In summary, an anionic phase-transfer protocol for enantioselective bromination was developed and applied to the 6-*exo*-trig bromocyclization of styrenyl amides. To realize this system, an insoluble cationic brominating reagent was developed, with optimal results delivered with 6,6'-silyl substituted TRIP phosphoric acid as an anionic phase-transfer precatalyst. Enantioselective iodination was achieved with an analogous cationic iodinating reagent. The resulting methods furnished a broad range of enantioenriched 4*H*-3,1-benzoxazines under mild and operationally simple reaction conditions. The highly insoluble nature of the halogenating reagents in nonpolar solvents effectively suppressed the noncatalyzed background reaction and led to halocyclization reactions with excellent enantioselectivity, even at low catalyst loading. For

example, reaction of cyclohexenyl substrate **3j** with **1d** with catalyst loading reduced of 0.1 mol % afforded the desired product in 87% yield and 90–93% ee.¹³ These methods extend the utility of the chiral anion phase-transfer approach and serve as a starting point for the design of other reagents for use in this paradigm.

■ ASSOCIATED CONTENT

■ Supporting Information

Experimental details, characterization data for new compounds, and crystallographic data (CIF). This material is available free of charge via the Internet at <http://pubs.acs.org>.

■ AUTHOR INFORMATION

Corresponding Author

fdtoste@berkeley.edu

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

We are grateful to the University of California Berkeley and to Amgen for financial support. V.R. thanks the NSERC for a postdoctoral fellowship. We gratefully acknowledge William J. Wolf for obtaining X-ray crystallographic data for **1d**.

■ REFERENCES

(1) Halogenation in natural products: (a) Wagner, C.; El Omari, M.; König, G. M. *J. Nat. Prod.* **2009**, *72*, 540. (b) Vaillancourt, F. H.; Yeh, E.; Vosburg, D. A.; Garneau-Tsodikova, S.; Walsh, C. T. *Chem. Rev.* **2006**, *106*, 3364. (c) Gribble, G. W. *J. Nat. Prod.* **1992**, *55*, 1353. For recent advances in bromocyclization in natural product synthesis, see: (d) Werness, J. B.; Tang, W. *Org. Lett.* **2011**, *13*, 3664. (e) Snyder, S. A.; Treiter, D. S. *Angew. Chem., Int. Ed.* **2009**, *49*, 7899. (f) Lin, H.; Pochapsky, S. S.; Krauss, I. J. *Org. Lett.* **2011**, *13*, 1222.

(2) Selected reviews on enantioselective halogenation: (a) Hennecke, U. *Chem. Asian J.* **2012**, *7*, 456. (b) Tan, C. K.; Zhou, L.; Yeung, Y.-Y. *Synlett* **2011**, 1335. (c) Castellanos, A.; Fletcher, S. P. *Chem.—Eur. J.* **2011**, *17*, 5766. (d) Ma, S.; Chen, G. *Angew. Chem., Int. Ed.* **2010**, *49*, 8306. (e) France, S.; Weatherwax, A.; Lectka, T. *Eur. J. Org. Chem.* **2005**, 475.

(3) Iodoetherification: (a) Kang, S. H.; Lee, S. B.; Park, C. M. *J. Am. Chem. Soc.* **2003**, *125*, 15748. Chlorohydrin synthesis: (b) Sakurada, I.; Yamasaki, S.; Göttlich, R.; Iida, T.; Kanai, M.; Shibasaki, M. *J. Am. Chem. Soc.* **2000**, *122*, 1245. (c) El-Qisairi, A.; Hamed, O.; Henry, P. M. *J. Org. Chem.* **1998**, *63*, 2790. Iodocarbocyclization: (d) Inoue, T.; Kitagawa, O.; Kurumizawa, S.; Ochiai, O.; Taguchi, T. *Tetrahedron Lett.* **1995**, *36*, 1479. Aminohalogenation: (e) Bovino, M. T.; Chemler, S. R. *Angew. Chem., Int. Ed.* **2012**, *51*, 3923.

(4) Halogenative semipinacol rearrangement: (a) Li, H.; Zhang, F.-M.; Tu, Y.-Q.; Zhang, Q.-W.; Chen, Z.-M.; Chen, Z.-H.; Li, J. *Chem. Sci.* **2011**, *2*, 1839. (b) Chen, Z.-M.; Zhang, Q.-W.; Zhang, Z.-H.; Li, H.; Tu, Y.-Q.; Zhang, F.-M.; Tan, J.-M. *J. Am. Chem. Soc.* **2011**, *133*, 8818. Dichlorination: (c) Nicolaou, K. C.; Simmons, N. L.; Ying, Y.; Heretsch, P. M.; Chen, J. S. *J. Am. Chem. Soc.* **2011**, *133*, 8134. (d) Juliá, S.; Ginebreda, A. *Tetrahedron Lett.* **1979**, *20*, 2171. Fluorocyclization: (e) Lozano, O.; Blessley, G.; del Campo, T. M.; Thompson, A. L.; Giuffredi, G. T.; Bettati, M.; Walker, M.; Borman, R.; Gouverneur, V. *Angew. Chem., Int. Ed.* **2011**, *50*, 8105. Chlorocyclization: (f) Jaganathan, A.; Garzan, A.; Whitehead, D. C.; Staples, R. J.; Borhan, B. *Angew. Chem., Int. Ed.* **2011**, *50*, 2593. Chlorolactonization: (g) Whitehead, D. C.; Yousefi, R.; Jaganathan, A.; Borhan, B. *J. Am. Chem. Soc.* **2010**, *132*, 3298. Bromolactonization: (h) Zhou, L.; Tan, C. K.; Jiang, X.; Chen, F.; Yeung, Y.-Y. *J. Am. Chem. Soc.* **2010**, *132*, 15474. (i) Zhang, W.; Zheng, S.; Liu, N.; Werness, J. B.; Guzei, I. A.; Tang, W. *J. Am. Chem. Soc.* **2010**, *132*, 3664. (j) Chen, J.; Zhou, L.; Tan, C. K.; Yeung, Y.-Y. *J. Org. Chem.* **2012**, *77*, 999.

(k) Zhang, W.; Liu, N.; Schienebeck, C. M.; Decloux, K.; Zheng, S.; Werness, J. B.; Tang, W. *Chem. Eur. — J.* **2012**, *18*, 7296. Bromoamination: (l) Zhou, L.; Chen, J.; Tan, C. K.; Yeung, Y.-Y. *J. Am. Chem. Soc.* **2011**, *133*, 9164. (m) Chen, J.; Zhou, L.; Yeung, Y.-Y. *Org. Biomol. Chem.* **2012**, *10*, 3808.

(5) Iodolactonization: Veitch, G. E.; Jacobsen, E. N. *Angew. Chem., Int. Ed.* **2010**, *49*, 7332.

(6) (a) Iodolactonization: Dodish, M. C.; Johnston, J. N. *J. Am. Chem. Soc.* **2012**, *134*, 6068. After submission of this work, the use of a BINOL-derived amidine-catalyzed bromolactonization was reported: (b) Paull, D. H.; Fang, C.; Donald, J. R.; Pansick, A. D.; Martin, S. F. *J. Am. Chem. Soc.* **2012**, *134*, 11129.

(7) Bromolactonization: (a) Murai, K.; Matsushita, T.; Nakamura, A.; Fukushima, S.; Shimura, M.; Fujioka, H. *Angew. Chem., Int. Ed.* **2010**, *49*, 9174. (b) Murai, K.; Nakamura, A.; Matsushita, T.; Shimura, M.; Fujioka, H. *Chem. Eur.—J.* **2012**, *18*, 8448.

(8) (a) Huang, D.; Wang, H.; Xue, F.; Guan, H.; Li, L.; Peng, X.; Shi, Y. *Org. Lett.* **2011**, *13*, 6350. (b) Hennecke, U.; Müller, C. H.; Fröhlich, R. *Org. Lett.* **2011**, *13*, 860. Use of a Lewis basic cocatalyst: (c) Denmark, S. E.; Burke, M. T. *Org. Lett.* **2012**, *14*, 256. Earlier study of Lewis base catalysis: (d) Denmark, S. A.; Burke, M. T. *Proc. Natl. Acad. Sci. U.S.A.* **2010**, *107*, 20655. (e) While this manuscript was under preparation, an enantioselective bromoamination of enamides with chiral phosphoric acids was reported: Alix, A.; Lalli, C.; Retailliau, P.; Masson, G. *J. Am. Chem. Soc.* **2012**, *134*, 10389.

(9) (a) Rauniyar, V.; Lackner, A. D.; Hamilton, G. L.; Toste, F. D. *Science* **2011**, *334*, 1681. (b) Phipps, R. J.; Hiramatsu, K.; Toste, F. D. *J. Am. Chem. Soc.* **2012**, *134*, 8376. See also: (c) Hamilton, G. L.; Kanai, T.; Toste, F. D. *J. Am. Chem. Soc.* **2008**, *130*, 14984. For a review see: (d) Phipps, R. J.; Hamilton, G. L.; Toste, F. D. *Nature Chem.* **2012**, *4*, 613.

(10) No enantioselective catalytic approaches to 4H-3,1-benzoxazines have been reported. For an elegant approach to the asymmetric synthesis of 4H-3,1-benzoxazin-2-ones in the context of HIV-1 reverse transcriptase inhibitor Efavirenz (Sustiva), see: Pierce, M. P.; Parsons, R. L., Jr.; Radesca, L. A.; Lo, Y. S.; Silverman, S.; Moore, J. R.; Islam, Q.; Choudhury, A.; Fortunak, J. M. D.; Nguyen, D.; Luo, C.; Morgan, S. J.; Davis, W. P.; Confalone, P. N.; Chen, C.; Tillyer, R. D.; Frey, L.; Tan, L.; Xu, F.; Zhao, D.; Thompson, A. S.; Corley, E. G.; Grabowski, E. J. J.; Reamer, R.; Reider, P. J. *J. Org. Chem.* **1998**, *63*, 8538. For an enantioselective catalytic synthesis of oxazines by 6-endo-trig chlorocyclization, see ref 4f.

(11) Early structural studies: (a) Blair, L. K.; Parris, K. D.; Hii, P. S.; Brock, C. P. *J. Am. Chem. Soc.* **1983**, *105*, 3649. (b) Bowmaker, G. A.; Hannan, S. F. *Aust. J. Chem.* **1972**, *25*, 1151. For mechanistic studies on the reaction of (bis)amino halonium reagents with alkenes see: (c) Neverov, A. A.; Brown, R. S. *J. Org. Chem.* **1998**, *63*, 5977. (d) Neverov, A. A.; Feng, H. X.; Hamilton, K.; Brown, R. S. *J. Org. Chem.* **2003**, *68*, 3802. (e) Brown, R. S.; Neverov, A. A.; Liu, C. T.; Maxwell, C. I. *ACS Symp. Ser.* **2007**, *965*, 458.

(12) (a) DABCO-bromine complex appears to be a complex oligomeric reagent whose structure depends on the exact conditions of synthesis. The sample used was prepared as in Heravi, M. M.; Derikvand, F.; Ghassemzadeh, M.; Neumüller, B. *Tetrahedron Lett.* **2005**, *46*, 6243. (b) Bis(*sym*-collidine)bromine(I) hexafluorophosphate was prepared as in Homsí, F.; Robin, S.; Rousseau, G. *Org. Synth.* **2000**, *77*, 206. (c) *N*-Bromosuccinimide was also screened, giving 37% and 27% ee with full conversion in 0.5 h, in the presence and absence of Na₂CO₃, respectively.

(13) Freshly dried and powdered base and vigorous stirring was required to achieve high enantioselectivities with 0.1 mol % catalyst loading. The reported range represents results obtained from 4 experiments.

■ NOTE ADDED AFTER ASAP PUBLICATION

This paper was published ASAP on July 25, 2012. Due to a production error, Table 2 was incorrect. The revised version was posted on July 26, 2012.